

CORRECTION

Open Access

Correction: phase I study of neoadjuvant chemoradiotherapy with S-1 plus biweekly cisplatin for advanced gastric cancer patients with lymph node metastasis -KOGC04-

Satoru Matsuda¹, Tsunehiro Takahashi^{1*}, Junichi Fukada², Kazumasa Fukuda¹, Hirofumi Kawakubo¹, Yoshiro Saikawa¹, Osamu Kawaguchi², Hiroya Takeuchi¹, Naoyuki Shigematsu² and Yuko Kitagawa¹

Correction

After the publication of this work [1], we noted that owing to an inadvertent mistake, the description of the chemotherapy schedule shown in the Methods was different from that shown in the Abstract and the duration of S-1 treatment in the initial chemoradiotherapy was inadequately described in Figure 1. Therefore, Methods and Figure 1 were modified accordingly.


In the “Chemoradiotherapy” section in the Methods, the 1st and 3rd paragraphs should read as follows:

“The chemoradiotherapy protocol consisted of administration of S-1 plus biweekly cisplatin and radiation (Figure 1). The initial chemoradiotherapy schedule was for 6 weeks: S-1 was orally administered twice daily from the evening of day 1 to the morning of day 15, and the total dose was based on the patient’s body surface area

(BSA), as follows: 1.25 m^2, 80 mg; $1.25\text{--}1.5\text{ m}^2$, 100 mg; and $>1.5\text{ m}^2$, 120 mg. An escalating dose of cisplatin was administered by infusion over 1 h on days 1 and 15 without infusional hydration. The starting dose (level 1) of cisplatin was 15 mg/m^2, the second dose (level 2) was 20 mg/m^2, and the third dose (level 3) was 25 mg/m^2.”

“After initial chemoradiotherapy, one cycle of combination chemotherapy with S-1 plus biweekly cisplatin was delivered. This consisted of 42 days of S-1 administered from the evening of day 1 to the morning of day 29 and of cisplatin administered on days 1, 15, and 29.”

In “Figure 1,” based on the description in the manuscript, the duration of S-1 treatment in the initial chemoradiotherapy was revised to 1–15 days from 1–22 days (Figure 1).



* Correspondence: surgeontsune@yahoo.co.jp

¹Department of Surgery, Keio University School of Medicine, 35 Shinanomachi, Shinjuku-ku, Tokyo 160-8582, Japan

Full list of author information is available at the end of the article

Author details

¹Department of Surgery, Keio University School of Medicine, 35 Shinanomachi, Shinjuku-ku, Tokyo 160-8582, Japan. ²Department of Radiology, Keio University School of Medicine, 35 Shinanomachi, Shinjuku-ku, Tokyo 160-8582, Japan.

Received: 16 June 2014 Accepted: 16 June 2014

Published: 25 June 2014

Reference

1. Matsuda S, Takahashi T, Fukada J, Fukuda K, Kawakubo H, Saikawa Y, Kawaguchi O, Takeuchi H, Shigematsu N, Kitagawa Y: **Phase I study of neoadjuvant chemoradiotherapy with S-1 plus biweekly cisplatin for advanced gastric cancer patients with lymph node metastasis: -KOGC04.** *Radiat Oncol* 2014, **9**:9.

doi:10.1186/1748-717X-9-140

Cite this article as: Matsuda *et al.*: Correction: phase I study of neoadjuvant chemoradiotherapy with S-1 plus biweekly cisplatin for advanced gastric cancer patients with lymph node metastasis -KOGC04-. *Radiation Oncology* 2014 **9**:140.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

